

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07K 14/54, A61K 38/20, C12N 15/24,</b> <b>G01N 33/74</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 96/18648</b> <b>(43) International Publication Date:</b> 20 June 1996 (20.06.96)
<b>(21) International Application Number:</b> PCT/IT95/00216 <b>(22) International Filing Date:</b> 13 December 1995 (13.12.95)  <b>(30) Priority Data:</b> RM94A000805 14 December 1994 (14.12.94) IT  <b>(71) Applicant (for all designated States except US):</b> ISTITUTO DI RICERCHE DI BIOLOGIA MOLECOLARE P. ANGELETTI S.P.A. [IT/IT]; Via Pontina Km. 30.600, I-00040 Pomezia (IT).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> CILIBERTO, Gennaro [IT/IT]; Viale Gorgia di Leontini, 330/19, I-00124 Casalpalocco (IT). SAVINO, Rocco [IT/IT]; Via della Tecnica, 76, I-00040 Roma (IT). LAHM, Armin [IT/IT]; Piazza S. Maria Ausiliatrice, 18, I-00153 Roma (IT). TONIATTI, Carlo [IT/IT]; Via Benedetto Croce, 26, I-00142 Roma (IT).  <b>(74) Agents:</b> DI CERBO, Mario et al.; Società Italiana Brevetti S.p.A., Piazza di Pietra, 39, I-00186 Roma (IT).		<b>(81) Designated States:</b> AU, CA, CN, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

**(54) Title:** SUPERAGONISTS AND ANTAGONISTS OF H IL-6, AND 3D MODELLING METHOD FOR THEIR SELECTION**(57) Abstract**

It is known that the ligands of the group of cytokines similar to Interleukin 6 (IL-6), that is Oncostatin M (OSM), Leukemia Inhibitory Factor (LIF), Ciliary Neurotrophic Factor (CNTF) and Interleukin 11 (IL-11), induce the formation of a receptor complex of which the membrane molecule gp 130 is a part. The present invention refers to a methodology for selecting superagonists, antagonists and superantagonists of human interleukin-6 comprising the following operations: comparing the amino acid sequence of bovine granulocyte colony stimulating factor (bG-CSF) with the sequence of said hormone; and on the basis of the above comparison, formulating a three-dimensional model of said hormone, which allows the identification of residues that form the site of interaction with the specific receptor (Site 1) and those that constitute the site of interaction with gp 130 (Site 2) respectively. The invention allows the identification of these sites in human interleukin-6 and the isolation of variants having, with respect to the wild type hormone, a greater affinity for the specific receptor (superagonists and superantagonists) or affinity for gp 130 reduced or abolished (antagonists and superantagonists). The figure shows a scheme illustrating the methodology applied to identify site 1 and site 2 in the case of human interleukin-6. The invention also describes the obtaining of specific superagonists and superantagonists of interleukin-6 and the use of superantagonists as low dose inhibitors of the growth of human myeloma cells dependent on wild type interleukin-6.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

**Superagonists and antagonists of h IL-6, and 3D modelling method for their selection****DESCRIPTION**

5           The present invention relates to a methodology for selecting superagonists, antagonists and superantagonists of human interleukin-6 (hereinafter referred to also as h IL-6 or IL-6) based on three-dimensional modelling.

10           As is known, WO 92/21029 to Genentec Inc. teaches a method for determination of agonists or antagonists of growth hormones and ligands with a similar structural conformation. The potential agonists and antagonists are put into contact with a receptor for the hormone and this causes formation of a ternary complex consisting of a  
15           molecule of the potential agonist or antagonist and two molecules of such receptor for the hormone to be agonized or antagonized. Dimerization of receptors induced by a ligand molecule allows to conclude that the ligand has two different interaction sites (site 1 and site 2), on  
20           which it is possible to operate using mutagenesis to generate agonists or antagonists.

          It is known that the ligands in the group of cytokines similar to Interleukin 6 (IL-6), that is Oncostatin M (OSM), Leukemia Inhibitory Factor (LIF),  
25           Ciliary Neurotrophic Factor (CNTF), and Interleukin 11 (IL-11), induce the formation of a receptor complex of which the membrane molecule gp 130 is a part. In this receptor complex the specific receptor for each of these cytokines and the membrane molecule gp 130 are always  
30           present as common elements. It is thus possible to formulate the hypothesis that site 1 and site 2 bind to two different molecules in this class of hormones: site 1 to the specific receptor and site 2 to gp 130. Identification of the two sites is made possible, as will  
35           be seen more clearly from the following, by construction of a three-dimensional model of the receptor complex based on the functional similarity between sequences of

the human growth hormone (hGH) receptor and sequences of the receptors for the hormones in question. Isolation of variants that, with respect to the wild type hormone, have a greater affinity for the specific receptor (superagonists or superantagonists) is obtained by  
5 construction of filamentous phage libraries, for example M13, carrying the hormone, both in the wild type and mutant version.

According to the invention, the difference between  
10 the three-dimensional model, for example of IL-6, adopted here and the one adopted in WO92/21029 leads to identify different residues in helix A and C as constituents of site 2. In fact, according to present invention, for the construction of the IL-6 model not the growth hormone,  
15 but the structure of a different cytokine was used as template.

Modelling of the human interleukin 6 molecule is performed as follows. It is known, from data available in scientific literature, that the amino acidic sequence  
20 of human interleukin 6 shows similarities with that of the granulocyte colony stimulating factor (G-CSF). The three-dimensional structure of bovine granulocyte colony stimulating factor (bG-CSF), determined using X-ray crystallography, was used as template to develop a three-  
25 dimensional model of human IL-6 from residue 16 to 184. Firstly, the amino acidic sequence of human IL-6 was aligned with that of bG-CSF. On the basis of the derived alignment, the amino acidic residues in the bG-CSF three-dimensional structure were replaced by the corresponding  
30 residues of human IL-6 using molecular modelling program in a computerized interactive graphic unit. In the positions in which alignment involves either deletions or insertions (which suggests a different local structure in the interleukin 6 molecule) adjustments were made by  
35 applying the options provided by the molecular modelling program.

This three-dimensional model of interleukin 6, based on the bG-CSF structure, has enabled the identification of the two sites of interaction between human interleukin 6 and its two receptors: the low affinity receptor gp 80 (site 1) and the high affinity signal transducer receptor gp 130 (site 2). The following procedure was used to identify the two sites. From sequence comparison it is known that all the members of the family of hematopoietic receptors are related to each other by the fact that they share a domain, known as the cytokine binding domain. This similarity of sequences also indicates a high probability of structural similarity in corresponding parts of the various receptors, including the two interleukin 6 receptors, gp 80 and gp 130. The observation that the cytokines that bind to these receptors all have (or are predicted to have) a similar structure, that is a four helix bundle, strongly supports the notion that the interaction between these cytokines and their receptors, by means of the cytokine binding domain, must be very similar - albeit not identical - in biologically active complexes.

Considering that the three-dimensional structure of one of these receptor complexes (the complex made by growth hormone and the extra-cellular domain of the dimeric receptor for the growth hormone, i.e. GHbp) has been determined by means of X-ray crystallography, our bG-CSF built model of human interleukin 6 allows us to identify the potential sites of interaction between interleukin 6 and its two receptors gp 80 (site 1) and gp 130 (site 2). This has been accomplished, according to the present invention, by constructing a structural model of gp 80 and gp 130 based on the coordinates furnished by the X-ray crystallographic structure of the growth hormone receptor, and by substituting in such complex the growth hormone with our bG-CSF built model of human interleukin-6 (see fig. 1).

As is known, interleukin 6 is a polypeptide of 184 amino acids which, as described, belongs to the class of helical cytokines. Interleukin 6 is a multi-functional cytokine produced by various cell types. It acts as a differentiation and growth factor on cells of various lineages, such as for example cells in the immune system, hepatocytes, kidney cells, hematopoietic stem cells, keratinocytes and neurones.

Production of superagonists of interleukin 6 would allow the use of therapeutic doses lower than those required with wild type interleukin 6 in the treatment of numerous serious diseases. In fact, interleukin 6 has important and promising applications in the treatment of breast cancer, leukemia, and infectious diseases or diseases connected with disorders of bone marrow progenitor cells.

In addition superagonists of IL-6 could be used in protocols for ex vivo expansion of hematopoietic progenitor cells both in bone marrow transplantation and gene therapy.

On the other hand the production of antagonists or superantagonists of human interleukin 6 would allow inhibition of interleukin 6 in numerous diseases characterized by its excessive production, such as chronic autoimmune diseases, myeloma/plasmacytoma, post-menopausal osteoporosis and cancer cachexia.

The methodology for the selection of superagonists, antagonists or superantagonists of interleukin-6, according to the present invention, comprises the following operations:

- comparing the amino acid sequence of bovine granulocyte colony stimulating factor (bG-CSF) with the sequence of said hormone; and
- on the basis of the above comparison, formulating a three dimensional model of said hormone, which allows the identification of residues that form the site of interaction with the specific receptor (Site 1)

and those that constitute the site of interaction with gp 130 (Site 2) respectively.

For selection of superagonists of interleukin 6, the methodology according to the present invention further comprises the following additional operations:

- production of a series of phage libraries containing mutations of the following wild type residues of interleukin 6 (present in the form of fusion product with filamentous phage proteins):

helix A:

Ser 22, Glu 23, Asp 26, Arg 30, Leu 33, Ser 37, Arg 40, Glu 42;

loop AB:

Ser 52, Ser 53, Ala 56, Leu 57, Glu 59, Asn 60, Leu 62, Leu 64, Pro 65, Lys 66,

Met 67, Ala 68, Glu 69, Lys 70, Asp 71, Phe 74, Gln 75, Ser 76;

helix D:

His 164, Leu 165, Arg 168, Ser 169, Lys 171, Glu 172, Phe 173, Gln 175, Ser 176, Ser 177, Leu 178, Arg 179, Ala 180, Leu 181, Arg 182, Gln 183, Met 184.

- selection, from the mixed population of phages belonging to each individual phage library and expressing interleukin 6 mutants, of that or those with an affinity for the specific receptor greater than that of wild type interleukin; and

- identification of the best receptor binder amino acid sequence or sequences by sequencing of the DNA extracted from the selected phage particles.

In this case, a series of phage libraries can be produced containing mutations of said wild type residues of interleukin 6 present as a fusion product with the M13 pIII protein.

The methodology for selecting antagonists of interleukin 6 according to the present invention comprises - along with the operations indicated above for

molecular modelling of the human IL-6 protein and its receptor chains - the following operations:

5       - mutagenesis of the residues identified to form part of the site of interaction with gp 130 (Arg 30, Tyr 31, Gly 35, Ser 37, Ala 38, Ser 118, Lys 120, Val 121, Gln 124, Phe 125, Gln 127, Lys 128 and Lys 129), using conventional molecular biology techniques;

10       - evaluation of biological activity and affinity for the specific interleukin 6 receptor of the mutants produced as above, in order to identify variants of interleukin 6 whose affinity for the specific receptor is normal and that show reduction or loss of the biological activity; and

15       - evaluation of the above variants of interleukin 6 as antagonists for the biological activity of wild type interleukin 6 on human cell lines.

20       In case of obtaining of superantagonists of interleukin 6 by combination of the variants of amino acid sequences responsible for antagonist activity, identified as above, with amino acid mutations responsible for an increased affinity of the specific receptor for interleukin 6.

25       In the methodology for obtaining antagonists or superantagonists of interleukin 6, the mutagenesis of the residues identified as above can be performed using a molecular biology technique chosen from the group comprising Polymerase Chain Reaction, Primer Extension, Oligonucleotide Directed Mutagenesis, and combinations thereof.

30       The present invention is not limited to the methodology for selection of superagonists, antagonists or superantagonists of interleukin 6. On the contrary, it extends to molecules obtainable by said methodology of selection, i.e. to: superagonists of h IL-6, with the exception of the molecule called IL-6 IRA and carrying  
35       the               following               three               substitutions  
Gln175Ile/Ser176Arg/Gln183Ala; antagonists of h IL-6,



with the exception of three molecules with the following substitutions:

Tyr31Asp/Gly35Phe/Ser118Arg/Val121Asp (DFRD)

Tyr31Asp/Gly35Phe/Ser118Phe/Val121Asp (DFFD)

5 Tyr31Asp/Gly35Phe/Ser118Leu/Val121Asp (DFLD);

and superantagonists of h IL-6, with the exception of the molecule called Sant1 and carrying the following seven substitutions: Tyr31Asp/Gly35Phe/Ser118Arg/Val121Asp/Gln175Ile/Ser176Arg/Gln183Ala.

10 Up to this point a general description of the subject of the present invention has been given. With the aid of the following examples a detailed description of specific embodiments of the invention will now be given, with the purpose of giving a better understanding  
15 of the objects, characteristics, advantages and methods of application thereof.

Figure 1 shows a scheme illustrating the methodology applied to identify site 1 and site 2 in the case of human interleukin 6.

20 Figure 2 shows the increase in potency of three superantagonists according to the invention, i.e. Sant 3, Sant 4 and Sant 5, over antagonist Tyr31Asp/Gly35Phe/Ser118Arg/Val121Asp (the one letter codes have been used in the figure), with the increase of concentration.

25 DEPOSITS

E.Coli K12 bacteria - transformed using the plasmid pHenΔhIL-6 containing, from the recognition site of the restriction enzyme SalI to that for the restriction enzyme NotI, a nucleotidic sequence coding for the amino  
30 acid sequence of wild type human interleukin 6 - have been deposited on 10/6/1993 with The National Collection of Industrial and Marine Bacteria Ltd. (NCIMB), Aberdeen, Scotland, UK, with access number NCIMB 40563.

#### Example 1

35 Application of the methodology according to the present invention for the selection of superagonists as

interleukin 6 by means of mutagenesis of amino acid residues in the AB loop

5 The strategy consists in construction of a hybrid gene containing all the region coding for hIL-6 (SEQ ID NO:1) followed by the last 157 amino acids of protein pIII of the phage M13 and preceded by the sequence Pel B, which vectors the synthesized protein to the periplasmic space.

10 This construct allows the obtaining of phagemid particles displaying on their surface correctly folded and biologically active human interleukin 6.

A phage library was constructed containing mutations of residues Asp 71, Phe 74, Gln 75 and Ser 76 of interleukin 6, starting from the variant IL-6 IRA (substitutions Gln175Ile/Ser176Arg/Gln183Ala) described 15 in WO95/00852 and having an affinity for the receptor approximately five times greater than that of wild type human interleukin 6, present in the form of fusion product with protein pIII of filamentous phage M13. The library was constructed using the Primer Extension 20 technique. The mutagenic oligonucleotide is IL-6 DFQS, a 95 nucleotides oligo, whose sequence is SEQ ID NO: 2. Primer IL-6 DFQS introduces degenerations into codons coding for the amino acids 71 (wild type Asp), 74 (wild type Phe), 75 (wild type Gln) and 76 (wild type Ser). 25 The oligonucleotide IL-6 AB primer, whose sequence is SEQ ID NO: 3, was used as primer for the Primer Extension reaction. The two oligonucleotides were annealed in vitro, and the annealed oligonucleotides were used as 30 substrate for a Primer Extension reaction. The double-stranded DNA fragment thus obtained was then digested and ligated into the plasmid pHenΔhIL-6 in order to replace the wild type sequence with the mutated ones. The ligation product was inserted in bacteria, yielding 35 roughly three million independent transformants. The transformed bacteria were infected with the M13K07 helper

bacteriophage to generate the phage library (a library of phasmids).

The library underwent selection by incubation with magnetic beads coated with monoclonal antibody directed against shrIL-6R and in the presence of shrIL-6R and shrgrp130. The phasmid population eluted at pH 3.6 was then amplified in bacteria. After four cycles of selection-amplification, randomly selected phasmids were sequenced over the mutagenized region, the corresponding mutant interleukin 6 proteins were produced in the periplasmic space of the appropriate bacterial strain and tested for interleukin 6 specific receptor binding. Table 1 shows that, using the methodology according to the present invention, it is possible to select variants of interleukin 6 having an additional increase in the affinity for the specific receptor, molecules with mutations both in helix D and in region A-B.

TABLE 1

Receptor binding properties in variants of interleukin 6-IRA containing additional mutations in the residues 71, 74, 75 and 76 of the region A-B

Position	71	74	75	76	Receptor binding (%)
wild type	Asp	Phe	Gln	Ser	100%
IL-6IRA	Asp	Phe	Gln	Ser	450%
phasmid D3-3	Asp	Tyr	Phe	Ile	2350%
phasmid D4-1	Asp	Tyr	Tyr	Val	2750%
phasmid D3-7	Asp	Phe	Tyr	Ile	2770%
phasmid D4-19	Asp	Phe	Tyr	Ser	1800%
phasmid D4-20	Asp	Phe	Tyr	Lys	4200%
phasmid D3-16	Asp	Phe	Tyr	Leu	1450%
phasmid D4-17	Asp	Phe	Phe	Ile	2430%

Example 2

Application of the methodology according to the present invention for obtaining superantagonists of interleukin 6

The four mutations Tyr31Asp/Gly35Phe/Ser118Arg/Vall21Asp (DFRD) confer antagonistic properties as described in WO95/00852. These four mutations were combined with mutations capable of increasing the specific receptor binding capacity (described in example 1), using the Polymerase Chain Reaction (PCR) molecular biology technique. More specifically:

the super-binder mutations on helix D and region AB of the phasmid D 3-7 (described in example 1), to create the mutant protein Sant 3;

the super-binder mutations on helix D and region AB of the phasmid D 3-3 (described in example 1), to create the mutant protein Sant 4;

the super-binder mutations on helix D and region AB of the phasmid D 4-20 (described in example 1), to create the mutant protein Sant 5.

The mutant proteins, containing nine (Sant 3 and Sant 5) or ten (Sant 4) amino acid substitutions, were tested both for their specific interleukin-6 receptor binding, and for their ability to antagonize the biological activity of interleukin-6 on human hepatoma and myeloma cells. Table 2 and fig. 2 show the specific receptor binding properties of DFRD and of Sant 3, Sant 4 and Sant 5 along with the concentrations (expressed in nanograms of mutant per milliliter of culture medium) of mutant necessary to inhibit 50% of interleukin 6 biological activity (hepatoma cells were stimulated with 4 nanograms of wild type interleukin 6 per milliliter of culture medium, while myeloma cells were stimulated with 0.1 nanograms of interleukin 6 per milliliter of culture medium, due to the higher sensitivity of the latter cells to wild type interleukin 6).

TABLE 2

Inhibition of wild type interleukin 6 biological activity on both human hepatoma and myeloma cells as a function of the mutant antagonists' specific interleukin-6 receptor binding capacity

---

Antagonist	Receptor binding	50% inhibition of interleukin 6 activity on:	
	(% of wild type)	hepatoma cells Hep3B	myeloma cells XG-1
5 DFRD	97%	164 ng/ml	190.0 ng/ml
Sant 3	2800%	2.4 ng/ml	1.85 ng/ml
Sant 4	2000%	2.7 ng/ml	3.90 ng/ml
Sant 5	4500%	2.3 ng/ml	2.45 ng/ml

---

- 10 As can be seen from the table, the introduction of the amino acid substitutions described in example 1 has at once increased the specific receptor binding capacity of the parental mutant DFRD and decreased the amount of antagonist needed to inhibit 50% of wild type interleukin
- 15 6 biological activity on both cell lines tested, therefore generating very effective and strong interleukin 6 superantagonists.

## SEQUENCE LISTING

## GENERAL INFORMATION

- (i) APPLICANT: ISTITUTO DI RICERCHE DI BIOLOGIA MOLECOLARE P. ANGELETTI S.p.A.
- 5 (ii) TITLE OF INVENTION: A METHODOLOGY FOR SELECTING SUPERAGONISTS, ANTAGONISTS AND SUPERANTAGONISTS OF HUMAN INTERLEUKIN-6 BASED ON RECEPTOR COMPLEX THREE DIMENSIONAL MODELLING
- (iii) NUMBER OF SEQUENCES: 3
- 10 (iv) CORRESPONDENCE ADDRESS:
- (A) ADDRESSEE: Società Italiana Brevetti
- (B) STREET: Piazza di Pietra, 39
- (C) CITY: Rome
- (D) COUNTRY: Italy
- 15 (E) POSTAL CODE: I-00186
- (v) COMPUTER READABLE FORM:
- (A) MEDIUM TYPE: Floppy disk 3.5" 1.44 MBYTES
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC DOS/MS DOS Rev 6.22
- 20 (D) SOFTWARE: Microsoft Word 6.0
- (viii) ATTORNEY INFORMATION
- (A) NAME: DI CERBO, Mario (Dr.)
- (B) REGISTRATION NUMBER:
- (C) REFERENCE: RM/X88471/PC-DC
- 25 (ix) TELECOMMUNICATION INFORMATION
- (A) TELEPHONE: 06/6785941
- (B) TELEFAX: 06/6794692
- (C) TELEX: 612287 ROPAT
- (1) INFORMATION FOR SEQ ID NO: 1:
- 30 (i) SEQUENCE CHARACTERISTICS
- (A) LENGTH: 555 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- 35 (ii) MOLECULE TYPE: DNA
- (iii) HYPOTHETICAL: no
- (iv) ANTISENSE: no

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE:

(A) SYNTHESIS: production in bacteria

(ix) FEATURE:

5 (A) NAME: IL-6 cDNA

(C) IDENTIFICATION METHOD: polyacrylamide

gel

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

	CCA	GTA	CCC	CCA	GGA	GAA	GAT	TCC	AAA	GAT	GTA	GCC	GCC	CCA	CAC	AGA	48
10	Pro	Val	Pro	Pro	Gly	Glu	Asp	Ser	Lys	Asp	Val	Ala	Ala	Pro	His	Arg	
	1				5					10					15		
	CAG	CCS	CTC	ACG	AGC	TCA	GAA	CGA	ATT	GAC	AAA	CAA	ATT	CGG	TAC	ATC	96
	Gln	Pro	Leu	Thr	Ser	Ser	Glu	Arg	Ile	Asp	Lys	Gln	Ile	Arg	Tyr	Ile	
				20					25					30			
15	CTC	GAC	GGC	ATC	TCA	GCC	TTA	AGA	AAG	GAG	ACA	TGT	AAC	AAG	AGT	AAC	144
	Leu	Asp	Gly	Ile	Ser	Ala	Leu	Arg	Lys	Glu	Thr	Cys	Asn	Lys	Ser	Asn	
			35					40					45				
	ATG	TGT	GAA	AGC	AGC	AAA	GAG	GCA	CTG	GCA	GAA	AAC	AAC	CTG	AAC	CTT	192
	Met	Cys	Glu	Ser	Ser	Lys	Glu	Ala	Leu	Ala	Glu	Asn	Asn	Leu	Asn	Leu	
20		50					55					60					
	CCA	AAG	ATG	GCT	GAA	AAA	GAT	GGA	TGC	TTC	CAA	TCT	GGA	TTC	AAT	GAG	240
	Pro	Lys	Met	Ala	Glu	Lys	Asp	Gly	Cys	Phe	Gln	Ser	Gly	Phe	Asn	Glu	
	65					70					75				80		
	GAG	ACT	TGC	CTG	GTG	AAA	ATC	ATC	ACT	GGT	CTT	TTG	GAG	TTT	GAG	GTA	288
25	Glu	Thr	Cys	Leu	Val	Lys	Ile	Ile	Thr	Gly	Leu	Leu	Glu	Phe	Glu	Val	
				85						90				95			
	TAC	CTA	GAG	TAC	CTC	CAG	AAC	AGA	TTT	GAG	AGT	AGT	GAG	GAA	CAA	GCC	336
	Tyr	Leu	Glu	Tyr	Leu	Gln	Asn	Arg	Phe	Glu	Ser	Ser	Glu	Glu	Gln	Ala	
				100					105					110			
30	AGA	GCT	GTC	CAG	ATG	AGT	ACA	AAA	GTC	CTG	ATC	CAG	TTC	CTG	CAG	AAA	384
	Arg	Ala	Val	Gln	Met	Ser	Thr	Lys	Val	Leu	Ile	Gln	Phe	Leu	Gln	Lys	
				115					120				125				
	AAG	GCA	AAG	AAT	CTA	GAT	GCA	ATA	ACC	ACC	CCT	GAC	CCA	ACC	ACA	AAT	432
	Lys	Ala	Lys	Asn	Leu	Asp	Ala	Ile	Thr	Thr	Pro	Asp	Pro	Thr	Thr	Asn	
35		130						135					140				

GCC AGC CTG CTG ACG AAG CTG CAG GCA CAG AAC CAG TGG CTG CAG GAC 480  
 Ala Ser Leu Leu Thr Lys Leu Gln Ala Gln Asn Gln Trp Leu Gln Asp  
 145 150 155 160  
 ATG ACA ACT CAT CTC ATT CTG AGA TCT TTT AAG GAG TTC CTG CAG TCC 528  
 5 Met Thr Thr His Leu Ile Leu Arg Ser Phe Lys Glu Phe Leu Gln Ser  
 165 170 175  
 AGC CTG AGG GCT CTT CGG CAA ATG TAG 555  
 Ser Leu Arg Ala Leu Arg Gln Met  
 180

10 (2) INFORMATION FOR SEQ ID NO: 2

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 95 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: synthetic DNA

(iii) HYPOTHETICAL: no

(iv) ANTISENSE: no

(v) FRAGMENT TYPE: internal

20

(vii) IMMEDIATE SOURCE:

(A) SYNTHESIS: oligonucleotide

synthesizer

(ix)

FEATURE:

(A) NAME: DFQS

(C) IDENTIFICATION METHOD: polyacrylamide

25

gel

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

GTGAGAGCTC CAAAGAGGCA CTGGCAGAAA ACAACCTGAA CCTTCCAAAG ATGGCTGAAA 60  
 AANNSSGGATG CNNSNNSNNS GGATTCAATG AGGAG 95

30

(3) INFORMATION FOR SEQ ID NO: 3

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 72 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: synthetic DNA

(iii) HYPOTHETICAL: no



(iv) ANTISENSE: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE:

(A) SYNTHESIS: oligonucleotide

5 synthesizer

(ix) FEATURE:

(A) NAME: AB primer

(C) IDENTIFICATION METHOD: polyacrylamide

gel

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GGCCTCTAGA TATACCTCAA ACTCCAAAAG ACCAGTGATG ATTTTCACCA GGCAAGTCTC 60  
CTCATTGAAT CC

72

## CLAIMS

1. A methodology for selecting superagonists, antagonists and superantagonists of interleukin 6 comprising the following operations:

- 5       - comparing the amino acid sequences of bovine granulocyte colony stimulating factor (bG-CSF) with the sequence of said hormone; and
- on the basis of the above comparison, formulating a three-dimensional model of said hormone, which allows the identification of residues that form the site of interaction with the specific receptor (site 1) and those that constitute the site of interaction with gp 130 (site 2) respectively.

15       2. The methodology for selecting superagonists of interleukin 6 according to claim 1, further comprising the following additional operations:

- production of a series of phage libraries containing mutations of the following wild type residues of interleukin 6 present in the form of a fusion product with filamentous phage proteins Ser 22, Glu 23, Asp 26, Arg 30, Leu 33, Ser 37, Arg 40, Glu 42, Ser 52, Ser 53, Ala 56, Leu 57, Glu 59, Asn 60, Leu 62, Leu 64, Pro 65, Lys 66, Met 67, Ala 68, Glu 69, Lys 70, Asp 71, Phe 74, Gln 75, Ser 76, His 164, Leu 165, Arg 168, Ser 169, Lys 171, Glu 172, Phe 173, Gln 175, Ser 176, Ser 177, Leu 178, Arg 179, Ala 180, Leu 181, Arg 182, Gln 183, Met 184;

- 30       - selection, from the mixed population of phages expressing interleukin 6 mutants, of that or those with an affinity for the specific receptor greater than that of wild type interleukin; and

      - identification of the best amino acid sequence or sequences binding the receptor by sequencing of the DNA extracted from the selected phage particles,

- 35       with the exception of the h IL-6 molecule carrying the three substitutions Gln175Ile/Ser176Arg/Gln183Ala.

3. The methodology for selecting superagonists of interleukin 6 according to claim 2, in which a series of phage libraries are produced, containing mutants of said wild type residue of interleukin 6 present as a product of fusion with protein pIII of M13.

4. The methodology for selecting antagonists of interleukin 6 according to claim 1, further comprising the following additional operations:

- mutation of the residues identified in claim 1, to form part of the site of interaction with gp 130 (Arg 30, Tyr 31, Gly 35, Ser 37, Ala 38, Ser 118, Lys 120, Val 121, Gln 124, Phe 125, Gln 127, Lys 128 and Lys 129), using conventional molecular biology techniques;

- evaluation of biological activity and affinity with the specific interleukin 6 receptor of the mutants produced as above, in order to identify variants of interleukin 6 whose affinity to the specific receptor is intact and that show reduction or loss of the biological activity; and

- evaluation of the above variants of interleukin 6 as antagonists for the biological activity of wild type interleukin 6,

with the exception of the three h IL-6 molecules with the following substitutions:

Tyr31Asp/Gly35Phe/Ser118Arg/Val121Asp  
Tyr31Asp/Gly35Phe/Ser118Phe/Val121Asp  
Tyr31Asp/Gly35Phe/Ser118Leu/Val121Asp.

5. The methodology for selecting superantagonists of interleukin 6 according to claims 2 to 4 by combination of the variations of amino acid sequences responsible for antagonist activity, indicated above, with amino acid variations responsible for an increased affinity of the specific receptor for interleukin 6, with the exception of the h IL-6 molecule carrying the seven substitutions  
Tyr31Asp/Gly35Phe/Ser118Arg/Val121Asp/Gln175Ile/Ser176Arg/Gln183Ala

6. The methodology for selecting antagonists or superantagonists of interleukin 6 according to claim 5, in which the mutagenesis of the residues identified as above is performed using a molecular biology technique chosen from the group comprising Polymerase Chain Reaction, Primer Extension, Oligonucleotide Directed Mutagenesis, and combinations thereof.

7. Interleukin 6 mutants according to claim 3, both showing an increased affinity for the specific receptor and containing the mutations Glutamine 175 Isoleucine, Serine 176 Arginine and Glutamine 183 Alanine, together with multiple substitutions in the Phenylalanine 74, Glutamine 75 and Serine 76 residues.

8. Interleukin-6 mutants according to claim 7, showing an increased affinity for the specific receptor and containing mutations chosen from the group comprising:

- Glutamine 75 Tyrosine, Serine 76 Isoleucine, Glutamine 175 Isoleucine, Serine 176 Arginine and Glutamine 183 Alanine;

- Phenylalanine 74 Tyrosine, Glutamine 75 Phenylalanine, Serine 76 Isoleucine, Glutamine 175 Isoleucine, Serine 176 Arginine and Glutamine 183 Alanine; and

- Glutamine 75 Tyrosine, Serine 76 Lysine, Glutamine 175 Isoleucine, Serine 176 Arginine and Glutamine 183 Alanine, said mutants exhibiting an affinity for the specific receptor increased 27.7 times, 23.5 times and 42 times, respectively.

9. Human interleukin-6 mutants having simultaneous substitution of the residues 31, 35, 74, 75, 76, 118, 121, 175, 176, 183 obtainable from claims 4 to 6 and which, by combining their antagonist properties to a greater affinity for the receptor, have the effect of superantagonists at low doses.

10. Human interleukin-6 mutants according to claim 9, with mutations chosen from the group comprising:

- Tyrosine 31 Aspartic acid, Glycine 35  
Phenylalanine, Serine 118 Arginine, Valine 121 Aspartic  
acid, Glutamine 75 Tyrosine, Serine 76 Isoleucine,  
Glutamine 175 Isoleucine, Serine 176 Arginine and  
5 Glutamine 183 Alanine;

- Tyrosine 31 Aspartic acid, Glycine 35  
Phenylalanine, Serine 118 Arginine, Valine 121 Aspartic  
acid, Phenylalanine 74 Tyrosine, Glutamine 75  
Phenylalanine, Serine 76 Isoleucine, Glutamine 175  
10 Isoleucine, Serine 176 Arginine and Glutamine 183  
Alanine; and


- Tyrosine 31 Aspartic acid, Glycine 35  
Phenylalanine, Serine 118 Arginine, Valine 121 Aspartic  
acid, Glutamine 75 Tyrosine, Serine 76 Lysine, Glutamine  
15 175 Isoleucine, Serine 176 Arginine and Glutamine 183  
Alanine,

said mutants being capable of inhibiting the  
biological activity of wild type interleukin 6 on  
sensitive human cells including myeloma cells the growth  
20 of which is IL-6-dependent.

11. Use of the superagonists according to claim 7 or  
8 for the preparation of drugs for therapy of  
trombocytopenia in man and for the ex vivo expansion of  
human hematopoietic progenitor cells for bone marrow  
25 transplantation and gene therapy.


12. Use of the interleukin-6 mutants according to  
claim 10 for the preparation of drugs for treatment of  
diseases characterized by overproduction of interleukin-  
6, and in particular of multiple myeloma, rheumatoid  
30 arthritis, postmenopausal osteoporosis and systemic lupus  
erythematosus.

# Construction of the Model<sup>1/2</sup>



## FIG. 1

**Increase of potency of Sant3, Sant4 and Sant5  
over antagonist Y31D/G35F/S118R/V121D**



**FIG. 2**

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/IT 95/00216

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C07K14/54 A61K38/20 C12N15/24 G01N33/74

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K A61K C12N G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EMBO JOURNAL, vol. 13, no. 6, 15 March 1994 pages 1357-1367, XP 000565719 SAVINO, R. ET AL. 'Generation of interleukin-6 receptor antagonists by molecular-modeling guided mutagenesis of residues important for gp 130 activation' * whole disclosure *	1-6
X	PROC. NATL. ACAD. SCI. USA, vol. 90, 1993 XP 000565720 SAVINO, R. ET AL. 'Saturation mutagenesis of the human interleukin 6 receptor binding site: Implications for its three-dimensional structure' * abstract; figs. 1-3 * --- -/-	1-3,5,6

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*A\* document member of the same patent family

Date of the actual completion of the international search

26 March 1996

Date of mailing of the international search report

12.04.96

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Hermann, R



# INTERNATIONAL SEARCH REPORT

Int. .onal Application No  
PCT/IT 95/00216

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO,A,94 09138 (CETUS ONCOLOGY CORPORATION) 28 April 1994 * claims 1-4 *	9,10,12
X	WO,A,94 11402 (ISTITUTO DI RICERCHE DI BIOLOGIA MOLECOLARE P. ANGELETTI S.P.A.) 26 May 1994 * page 1; claims *	1
A	J. IMMUNOL. , vol. 153, no. 4, 15 August 1994 pages 1744-1753, XP 000565715 EHLERS, M. ET AL. 'Identification of two novel regions of human IL-6 responsible for receptor binding and signal transduction' * abstract; p. 1746, left-hand column, last paragraph; p. 1749-1751 *	1-12

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Application No  
PCT/IT 95/00216

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
W0-A-9409138	28-04-94	AU-B-	5409294	09-05-94
		CA-A-	2147466	28-04-94
		EP-A-	0672144	20-09-95
-----				
W0-A-9411402	26-05-94	CA-A-	2129824	26-05-94
		EP-A-	0667872	23-08-95
		JP-T-	7501947	02-03-95
-----				